

Listeria monocytogenes*, Strain FSL J2-003*Catalog No. NR-13234****For research only. Not for human use.****Contributor:**

Professor Patrick L. McDonough, Ph.D., Director, Bacteriology and Mycology Section, Animal Health Diagnostic Center, Department of Population Medicine and Diagnostic Sciences, College of Veterinary Medicine, Cornell University, Ithaca, New York

Product Description:

Bacteria Classification: *Listeriaceae*, *Listeria*

Species: *Listeria monocytogenes*

Strain: FSL J2-303

Serotype: 1/2a

Original Source: *Listeria monocytogenes* (*L. monocytogenes*), strain FSL J2-003 was isolated from an aborted bovine fetus in New York, 1993.¹

Comment: The complete genome of *L. monocytogenes*, strain FSL J2-003 has been sequenced (GenBank: AARM00000000).² For more sequencing information, refer to the Broad Institute's [Listeria Genome Project](#).

L. monocytogenes is a Gram-positive, facultative intracellular bacterium that is extremely tolerant of external stresses (pH 3 to 12, temperatures ranging from 1 to 45°C, and high salt). *L. monocytogenes* encompasses a diversity of strains with varied virulence and pathogenic potential. There are 13 serotypes (1/2a, 1/2b, 1/2c, 3a, 3b, 3c, 4a, 4b, 4c, 4d, 4e, 5 and 7) that have been isolated from mammalian, bird, fish and shellfish species as well as environmental sources. Of these, only 3 serotypes (1/2a, 1/2b, and 4b) are frequently isolated from outbreaks of human listeriosis. The most common cause of infection is through ingestion of contaminated foods, in particular milk, meat or vegetable products. The infective dose is unknown and varies with species.³

L. monocytogenes, strain FSL J2-003 is a serotype 1/2a strain of ribotype DUP-1039C, one of the most common subtypes associated with ruminant listeriosis cases. It is also commonly isolated in feces of healthy ruminants and the farm environment.⁴

Material Provided:

Each vial contains approximately 0.5 mL of bacterial culture in 0.5X Brain Heart Infusion broth supplemented with 10% glycerol.

Note: If homogeneity is required for your intended use, please purify prior to initiating work.

Packaging/Storage:

NR-13234 was packaged aseptically, in screw-capped

plastic cryovials. The product is provided frozen and should be stored at -60°C or colder immediately upon arrival. For long-term storage, the vapor phase of a liquid nitrogen freezer is recommended. Freeze-thaw cycles should be avoided.

Growth Condition:**Media:**

Brain Heart Infusion broth or equivalent
Brain Heart Infusion agar or Sheep Blood agar

Incubation:

Temperature: 37°C
Atmosphere: Aerobic

Propagation:

1. Keep vial frozen until ready for use, then thaw.
2. Transfer the entire thawed aliquot into a single tube of broth.
3. Use several drops of the suspension to inoculate an agar slant and/or plate.
4. Incubate the tubes and plate at 37°C for 24 hours.

Citation:

Acknowledgment for publications should read "The following reagent was obtained through the NIH Biodefense and Emerging Infections Research Resources Repository, NIAID, NIH: *Listeria monocytogenes*, Strain FSL J2-003, NR-13234."

Biosafety Level: 2

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the following publication: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, and National Institutes of Health. [Biosafety in Microbiological and Biomedical Laboratories](#). 5th ed. Washington, DC: U.S. Government Printing Office, 2007; see www.cdc.gov/od/ohs/biosfty/bmbl5/bmbl5toc.htm.

Disclaimers:

You are authorized to use this product for research use only. It is not intended for human use.

Use of this product is subject to the terms and conditions of the BEI Resources Material Transfer Agreement (MTA). The MTA is available on our Web site at www.beiresources.org.

While BEI Resources uses reasonable efforts to include accurate and up-to-date information on this product sheet, neither ATCC® nor the U.S. Government make any warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. Neither ATCC® nor the U.S. Government warrants that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, use and disposal. ATCC® and the U.S. Government are not liable for any

SUPPORTING INFECTIOUS DISEASE RESEARCH

damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, the U.S. Government, ATCC®, their suppliers and contributors to BEI Resources are not liable for damages arising from the misidentification or misrepresentation of products.

Use Restrictions:

This material is distributed for internal research, non-commercial purposes only. This material, its product or its derivatives may not be distributed to third parties. Except as performed under a U.S. Government contract, individuals contemplating commercial use of the material, its products or its derivatives must contact the contributor to determine if a license is required. U.S. Government contractors may need a license before first commercial sale.

References:

1. Dr. Patrick McDonough, personal communication.
2. Glaser, P., et al. "Comparative Genomics of *Listeria* Species." Science 294 (2001): 849-852. PubMed: 11679669. GenBank: AARM00000000.
3. Liu, D., et al. "Toward an Improved Laboratory Definition of *Listeria monocytogenes* Virulence." Int. J. Food Microbiol. 118 (2007): 101-115. PubMed: 17727992.
4. Nightingale, K., et al. "Ecology and Transmission of *Listeria monocytogenes* Infecting Ruminants and in the Farm Environment." Appl. Environ. Microbiol. 70 (2004): 4458-4467.
5. Gray, M.J., et al. "*Listeria monocytogenes* Isolates from Foods and Humans Form Distinct but Overlapping Populations." Appl. Environ. Microbiol. 70 (2004): 5833-5841. PubMed: 15466521.
6. Boneca, I. G., et al. "A Critical Role for Peptidoglycan *N*-Deacetylation in *Listeria* Evasion from the Host Innate Immune System." Proc. Natl. Acad. Sci. U.S.A. 104 (2007): 997-1002. PubMed: 17215377.
7. Cai, S., et al. "Rational Design of DNA Sequence-Based Strategies for Subtyping *Listeria monocytogenes*." J. Clin. Microbiol. 40 (2002): 3319-3325. PubMed: 12202573.
8. Hain, T., C. Steinweg, and T. Chakraborty. "Comparative and Functional Genomics of *Listeria* Spp." J. Biotechnol. 126 (2006): 37-51. PubMed: 16757050.
9. Liu, D., et al. "Characterization of Virulent and Avirulent *Listeria monocytogenes* Strains by PCR Amplification of Putative Transcriptional Regulator and Internalin Genes." J. Med. Microbiol. 52 (2003): 1065-1070. PubMed: 14614064.
10. Milillo, S. R., J. M. Badamo and M. Wiedmann. "Contributions to Selected Phenotypic Characteristics of Large Species- and Lineage-specific Genomic Regions in *Listeria monocytogenes*." Food Microbiol. 26 (2009): 212-223. PubMed: 19171265.
11. Rocourt, J. and C. Buchrieser. "The Genus *Listeria* and *Listeria monocytogenes*: Phylogenetic Position, Taxonomy, and Identification." In: E. T. Ryser and E. H. Marth, Listeria, Listeriosis, and Food Safety (3rd ed.) New York: Marcel Dekker, Inc., pp 1-20.
12. Seeliger, H. P. R. Listeriosis. 2nd ed. Basel: Karger, 1961.

ATCC® is a trademark of the American Type Culture Collection.

