

Human Respiratory Syncytial Virus, A1998/3-2, Purified from HEp-2 Cells

Catalog No. NR-44233

For research use only. Not for use in humans.

Contributor:

Martin L. Moore, Ph. D., Department of Pediatrics, Emory University, Atlanta, Georgia, USA

Manufacturer:

BEI Resources

Product Description:

Virus Classification: *Pneumoviridae*, *Orthopneumovirus*, *Human orthopneumovirus*

Species: Human respiratory syncytial virus

Strain/Isolate: A1998/3-2

Original Source: Human respiratory syncytial virus (RSV), A1998/3-2 was isolated from a nasal wash from an infant with RSV bronchiolitis in Nashville, Tennessee, USA, on March 2, 1998.¹

Comments: A1998/3-2 is one of six clinical RSV isolates that recently were shown to induce variable disease severity, lung interleukin-13 (IL-13) levels, and gob-5 levels in BALB/cJ mice.² IL-13 is a cytokine linked to mucus production and gob-5 is a calcium-activated chloride channel family member implicated in airway inflammation.^{3,4} Compared to mock infection, RSV A1998/3-2 infection led to low levels of gob-5 in lung tissue, no significant elevation in IL-13 expression, and no weight loss in infected mice.²

NR-44233 was prepared by inoculation of human epithelial carcinoma cells (HEp-2; ATCC® CCL-23™) with RSV, A1998/3-2. The virus was purified from clarified supernatant by high speed centrifugation.

A similarly processed preparation of mock-infected HEp-2 cell clarified supernatant, suitable for use as a control, is available as BEI Resources NR-43974.

Material Provided:

Each vial contains approximately 0.5 mL of purified virus in TBS (0.15 M sodium chloride, 0.05 M Tris-HCl, pH 7.6).

Packaging/Storage:

NR-44233 was packaged aseptically in screw-capped plastic cryovials. The product is provided frozen and should be stored at -60°C or colder immediately upon arrival. For long-term storage, the vapor phase of a liquid nitrogen freezer is recommended. Freeze-thaw cycles should be avoided.

Growth Conditions:

Host: Human Epithelial carcinoma cells (HEp-2; ATCC® CCL-23™)

Growth Medium: Eagle's Minimum Essential Medium containing Earle's Balanced Salt Solution, non-essential

amino acids, 2 mM L-glutamine, 1 mM sodium pyruvate and 1.5 g/L of sodium bicarbonate supplemented with 2% fetal bovine serum, or equivalent

Infection: Cells should be 60% to 80% confluent

Incubation: 5 to 10 days at 37°C and 5% CO₂

Cytopathic Effect: Syncytia, cell rounding and sloughing

Citation:

Acknowledgment for publications should read "The following reagent was obtained through BEI Resources, NIAID, NIH: Human Respiratory Syncytial Virus, A1998/3-2, Purified from HEp-2 Cells, NR-44233."

Biosafety Level: 2

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the following publication: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, and National Institutes of Health. [Biosafety in Microbiological and Biomedical Laboratories \(BMBL\)](#), 6th ed. Washington, DC: U.S. Government Printing Office, 2020.

Disclaimers:

You are authorized to use this product for research use only. It is not intended for human use.

Use of this product is subject to the terms and conditions of the BEI Resources Material Transfer Agreement (MTA). The MTA is available on our Web site at www.beiresources.org.

While BEI Resources uses reasonable efforts to include accurate and up-to-date information on this product sheet, neither ATCC® nor the U.S. Government makes any warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. Neither ATCC® nor the U.S. Government warrants that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, use and disposal. ATCC® and the U.S. Government are not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, the U.S. Government, ATCC®, their suppliers and contributors to BEI Resources are not liable for damages arising from the misidentification or misrepresentation of products.

Use Restrictions:

This material is distributed for internal research, non-commercial purposes only. This material, its product or its derivatives may not be distributed to third parties. Except as performed under a U.S. Government contract, individuals contemplating commercial use of the material, its products or its derivatives must contact the contributor to determine if a license is required. U.S. Government contractors may need a license before first commercial sale.

References:

1. Moore, M. L., Personal Communication.
2. Stokes, K. L., et al. "Differential Pathogenesis of Respiratory Syncytial Virus Clinical Isolates in BALB/c Mice." *J. Virol.* 85 (2011): 5782-5793. PubMed: 21471228.
3. Nakanishi, A., et al. "Role of gob-5 in Mucus Overproduction and Airway Hyperresponsiveness in Asthma." *Proc. Natl. Acad. Sci. USA* 98 (2001): 5175-5180. PubMed: 11296262.
4. Walter, D. M., et al. "Critical Role for IL-13 in the Development of Allergen-Induced Airway Hyperreactivity." *J. Immunol.* 167 (2001): 4668-4675. PubMed: 11591797.

ATCC® is a trademark of the American Type Culture Collection.

