

Chikungunya Virus, YO 123223

Catalog No. NR-49908

Product Description: Cell lysate and supernatant from *Cercopithecus aethiops* kidney epithelial cells¹ infected with chikungunya virus (CHIKV), YO 123223

Passage History: C2V1/V2 (Prior to deposit at BEI Resources/BEI Resources); C = Number of passages in *Aedes albopictus* C6/36 cells; V# = Number of passages in Vero cells

Lot²: 63965678

Manufacturing Date: 26FEB2016

TEST	SPECIFICATIONS	RESULTS
Infectivity in Vero E6 Cells	Report results	Cell rounding and detachment
Sequencing of Species-Specific Region (914 nucleotides)	Consistent with CHIKV	Consistent with CHIKV ³
Titer by TCID₅₀ Assay^{4,5} in Vero E6 Cells¹	Report results	2.8 × 10 ⁶ TCID ₅₀ per mL
Amplification of CHIKV Sequence by RT-PCR	~ 1070 bp amplicon	~ 1070 bp amplicon
Sterility (21-day incubation) Harpo's HTYE broth ⁶ , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO ₂	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) DNA detection by PCR of extracted Test Article nucleic acid	None detected None detected	None detected None detected

¹Vero 76, clone E6; ATCC® CRL-1586™

²Grown in Eagle's Minimum Essential Medium containing Earle's Balanced Salt Solution, non-essential amino acids, 2 mM L-glutamine, 1 mM sodium pyruvate and 1.5 g/L of sodium bicarbonate (ATCC® 30-2003) supplemented with 2% fetal bovine serum (ATCC® 30-2020) for 3 days at 37°C with 5% CO₂

³Sequence information for CHIKV, YO 123223 is not available in the NCBI database; nucleotide sequence obtained for NR-49908, Lot No. 63965678 is ~ 99% identical to numerous CHIKV strains isolated in the Caribbean.


⁴The Tissue Culture Infectious Dose 50% (TCID₅₀) endpoint is the 50% infectious endpoint in cell culture. The TCID₅₀ is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID₅₀ provides a measure of the titer (or infectivity) of a virus preparation.

⁵5 days at 37°C and 5% CO₂

⁶Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

Date: 08 AUG 2016

Signature:



BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

